

## Canine Filariosis Around Istanbul, Turkey Employing Naphtol AS-TR Phosphatase

M. TOPARLAK, A. GARGILI, M. ULUTAS ESATGIL, H. CETINKAYA

Istanbul University, Veterinary Faculty, Department of Parasitology 34851,  
Istanbul, Turkey

Received April 15, 2003

Accepted June 6, 2005

### Abstract

Toparlak M., A. Gargili, M. Ulutas Esatgil, H. Cetinkaya: Canine Filariosis Around Istanbul, Turkey Employing Naphtol AS-TR Phosphatase technique. Acta Vet. Brno 2005, 74: 233-236.

The aim of this study was to find the prevalence of filarial nematodes in dogs in Istanbul, Turkey and identify the species by using naphtol AS-TR phosphatase technique. A total of 286 blood samples were taken from the dogs of different areas of Istanbul, Turkey, between March 1999 and November 2002. The Knott technique was used to detect the microfilariae in blood. Two dogs were found to be positive for microfilariae. The positive blood samples were stained by naphtol AS-TR phosphatase for differentiation. The microfilariae were identified as *Dipetalonema reconditum*. In this study, differentiation of canine microfilaria based on somatic distribution of acid phosphatase activity was performed for the first time and according to these results *D. reconditum* was reported for the first time from Turkey.

*Dog, Filaria, Dipetalonema reconditum, Knott-technique, Naphtol AS-TR Phosphatase*

Canine filariosis is a nematode disease caused by *Dirofilaria* spp., *Dipetalonema* spp. and *Brugia* spp. in dogs. Among the species *D. immitis* has a specific importance based on heart localization of adults and it infects also a wide variety of mammals including humans. The differentiation of microfilariae based on morphological peculiarities is difficult and naphtol AS-TR phosphatase technique is the most reliable method for their differentiation. Prevalence of dog filariae has been studied in various countries, in Turkey, no survey on prevalence of filarial nematodes of dog has been encountered except case reports based on finding of adults and morphological identification of microfilariae.

There are many species of filariae in dogs (Table 1). Three of them, *Dirofilaria immitis*, *D. repens* (Table 2) and *Dipetalonema reconditum* (Erdil 1966) have been reported from dogs in various parts of Turkey.

Although the most important host is the dog, *D. immitis* infects also a wide variety of mammals including human beings (Cheng 1986; Soulsby 1982). In this respect, the species identification of microfilariae is gaining in importance. It is reported that the differentiation of microfilariae based on morphological peculiarities is difficult and naphtol AS-TR technique is the most reliable method for their differentiation. It is based on acid phosphatase enzyme activity in the different body parts of each species of microfilariae (Acevedo et al. 1981; Chalifoux and Hunt 1971; Ortega-Mora et al. 1989; Schrey 1996; Whitlock et al. 1978). In this study, identification of filariid species in dogs was performed and the prevalence was determined in Istanbul for the first time.

### Materials and Methods

This survey was performed between March 1999 and November 2002. A total of 286 blood samples of stray dogs in the various parts of Istanbul were examined for filarial nematodes. All dogs were over 1 year-old and both sexes. About 5 ml blood was drawn from the cephalic vein of each dog (between 10:00 am-05:00 pm) and put into

#### Address for correspondence:

Dr. Aysen Gargili  
Istanbul University Veterinary Faculty  
Department of Parasitology  
34851, Avcilar, Istanbul, Turkey

Tel: 90 212 591 69 79  
Fax: 90 212 59169 91  
90 212 591 69 76  
e-mail: agargili@istanbul.edu.tr  
<http://www.vfu.cz/acta-vet/actavet.htm>

Table 1. Filarioid nematodes in dogs (Kelly 1979)

Species	Host Range	Vector	Tissue sites (adults)	Tissue sites (microfilariae)	Length of microfilariae (µm)	Geographical distribution
<i>Dirofilaria immitis</i>	Dog, cat, marine mammals, man	Mosquito	Heart and pulmonary artery	Blood	286-349 (mean:314)	America, Africa, Australia, Italy, Spain
<i>Dirofilaria repens</i>	Dog, cat, man	Mosquito	Subcutaneous tissues	Blood	200	USSR, Europe, India, Far East
<i>Dirofilaria conjunctivae</i>	Considered identical to <i>D. repens</i> and may also be confused with <i>D. tenuis</i> , which is found in the subcutaneous tissues of raccoons and man					
<i>Dipetalonema reconditum</i>	Dog	Fleas, ticks	Connective tissues	Blood	258-292 (mean:270)	America, Africa, Australia, Italy
<i>Dipetalonema dracunculoides</i>	Dog	Louse Fly ( <i>Hippobosca longifennis</i> )	Peritoneal membranes	Blood	195-230	Africa
<i>Dipetalonema grassi</i>	Dog	Tick	Subcutaneous tissues	Skin, rarely in blood	570	Italy, Kenya
<i>Brugia malayi</i>	Man, cat (dog)	Mosquito	Lymphatic system	Blood	170-260 (mean:220)	India, Africa, Far East
<i>Brugia pahangi</i>	Dog, cat, felidae	Mosquito	Lymphatic system	Blood	280	Africa, Far East
<i>Brugia patei</i>	Dog, cat	Mosquito	Lymphatic system	Blood	Similar to B.malayi	Africa

Table 2. Filarial nematodes of dogs in Turkey

Cities -Years	No of examined dogs	No of infected dogs and species found	Reference
Elazg-1983	283 (Knott Technique)	20 (7.07%) with <i>D. repens</i> -microfilaria 5 (1.77%) with <i>D. repens</i> +nonspecific-microfilaria 28 (9.89%) with nonspecific-microfilaria	Tasan 1983
Ankara-1983	50 (Necropsy)	2 (4%) with <i>D. repens</i> -adult	Doganay 1983
Elazg-1984	120 (Knott Technique)	1 (0.83%) with <i>D. immitis</i> -microfilaria 3 (2.5%) with <i>D. repens</i> -microfilaria 6 with <i>D. immitis</i> -adult 3 with <i>D. repens</i> -adult	Tasan 1984
Eskisehir-1986	2 (Necropsy) 20 (Knott Technique)	2 with <i>D. immitis</i> adult 6 with <i>D. immitis</i> microfilaria	Sarnic 1986
Bursa-1989	100 (Necropsy)	2 (2%) with <i>D. immitis</i> -adult	Tinar et al. 1989
Ankara-1989	27 (Necropsy)	3 (11.1%) with <i>D. immitis</i> -adult	Zeybek 1989
Konya-1990	4 (Necropsy)	4 with <i>D. immitis</i> -adult	Cantoray et al. 1990
Gemlik-1992	168 (Knott Technique)	5 (2.98%) with <i>D. immitis</i> -microfilaria	Coskun S.Z. et al. 1982
Van-1992	9 (Knott Technique) 10 (Necropsy)	7 with <i>D. immitis</i> -microfilaria 8 with <i>D. immitis</i> -adult	Agaoglu and Sahin 1992
Ankara-1992	33 (Necropsy)	3 (9.09%) with <i>D. immitis</i> -adult	Zeybek et al. 1992
Kayseri-1993	50 (Necropsy)	8 (16%) with <i>D. immitis</i> -adult	Sahin et al. 1993
Sivas-1997	50 (Necropsy)	3 (6%) with <i>D. immitis</i> -adult	Atas et al. 1997

a tube coated with lithium-heparin to prevent coagulation. Knott technique ( Sloss et al. 1994) was used for the detection of microfilariae. When microfilariae were seen, 2 ml of this positive sample was mixed with 12 ml of distilled water to lyse the red blood cells (Rossi and Abbate 1990). This mixture was centrifuged at 1500 rpm for 5 min. Supernatant was discarded and thin smears were prepared from the sediment. Smears were air dried, fixed in absolute acetone for 1-2 min at 4 °C and stained with naphtol AS-TR phosphatase (Sigma, St. Louis, USA, Catalog No. N6125) technique (Chalifoux and Hunt 1971). Identification of the microfilariae was done according to Acevedo et al. (1981), Chalifoux and Hunt (1971) and Schrey (1996).

## Results

Two of the 286 (0.69%) examined dogs had microfilariae in their blood. Only one type of microfilariae were observed in smears stained with naphtol AS-TR phosphatase technique.

The entire body of each microfilaria was stained bright red and higher enzyme activity was noticed between the excretory and anal pore. These microfilariae were identified to be *Dipetalonema reconditum*.

The dimension of microfilariae varied between  $260\text{-}290 \times 4.5\text{-}6 \mu\text{m}$  (average:  $275 \times 5.25 \mu\text{m}$ ). The districts and the numbers of examined and infected dogs are given in Table 3.

Table 3. The districts, the numbers of examined and infected dogs and the species found

Districts	No. of examined dogs	No. of infected dogs	Infection rate (%)	Species found
Büyükecemece	51			-
Haramidere	20			-
Kadıköy	39			-
Ümraniye	44	1	2.27	<i>Dip. reconditum</i>
Kemerburgaz	50			-
Zeytinburnu	14			-
Alibeyköy	18			-
Sarıyer	50	1	2	<i>Dip. reconditum</i>
Total	286	2	0.7	

### Discussion

Among many filarial nematodes in dogs (Kelly 1979; Soulsby 1982) mainly 2 species, namely *Dirofilaria immitis* and *D. repens* occur in Turkey (Table 2). Although the presence of *D. reconditum* has been reported from one dog, results were based only on the length, width and tail morphology of the microfilariae and therefore suspected (Erdil 1966). As it is seen in Table 1, other species are distributed mostly Africa, India and Far East. Adults of canine filariae inhabit the subcutaneous tissue or lymphatic system, only *D. immitis* adults reside inside the heart. Because of the ability of this parasite to infect the human beings in the same pattern, it is the most dangerous species among the dog filariae and its possible public health importance has been discussed in the studies (Atas et al. 1997; Doganay 1983) reporting *D. immitis* and conducted in other parts of the Turkey. *D. repens* and *D. reconditum* are considered to be non-pathogenic (Acevedo et al. 1981; Kelly 1979; Lindsey 1965; Soulsby 1982). Therefore it is very important to make an accurate differential diagnosis of microfilariae of these non-pathogenic species from the pathogenic and zoonotic *D. immitis*. In the previous studies (Agaoglu and Sahin 1992; Coskun et al. 1992; Doganay 1983; Sarnic and Alkan 1986; Tasan 1983; Tasan 1984) performed in Turkey, morphological features and length-width measurements were used for the identification of microfilariae. No doubt that this procedure is time-consuming and difficult whatever technique is employed.

The present findings show that the only species found in dogs in Istanbul was *D. reconditum* and naphthol AS-TR phosphatase technique based on acid phosphatase activity given an accurate diagnostic criteria since the microfilariae of 3 species of dogs have been reported (Acevedo et al. 1981; Balbo and Abate 1972; Chalifoux and Hunt 1971; Whitlock et al. 1978) to have completely different staining pattern. In this study the staining pattern of *D. reconditum* fits the original description by Chalifoux and Hunt (1971). According to the results of the present study, the major filaria of the dogs in Istanbul is *D. reconditum*. Although no pathogenic effects have been ascribed to this parasite and also it is not of public health importance compared to *D. immitis*, it is important to accurately diagnose and differentiate its microfilariae from those of *D. immitis*, in the diagnosis of heartworm infections.

### Výzkum filariózy u psů v okolí Istanbulu pomocí naftol AS-TR fosfatázové techniky

Cílem studie bylo zjistit prevalenci filárií u psů v okolí Istanbulu a určit jejich druh pomocí naftol AS-TR fosfatázy. Od března 1999 do listopadu 2002 bylo odebráno celkem 286 vzorků krve psů z různých oblastí Istanbulu. K detekci mikrofilárií v krvi byla použita

Knottova metoda. Test byl pozitivní u dvou psů. Pozitivní krevní vzorky byly k diferenciaci obarveny naftol AS-TR fosfatázou. Mikrofilárie byly identifikovány jako *Dipetalonema reconditum*. V této studii byly poprvé v Turecku mikrofilárie diferencovány na základě somatické distribuce aktivity kyselých fosfatáz, identifikovány jako *D. reconditum*.

#### References

- ACEVEDO RA, THEIS JH, KRAUS JF, LONGHURST WM 1981: Combination of filtration and histochemical stain for detection and differentiation of *Dirofilaria immitis* and *Dipetalonema reconditum* in the dog. *Am J Vet Res* **42**: 537-540
- AGAOGLU ZT, SAHIN A 1992: Van'da *Dirofilaria immitis* (*Dirofilaria immitis* in Van). *YY Üniv Vet Fak Derg* **3**(1-2): 117-121
- ATAS AD, ÖZCELİK S, SAYGI G 1997: Sivas sokak köpeklerinde görülen helmint türleri, bunların yayılışı ve halk sağlığı yönünden önemi (The occurrence of Helminth Species in Stray Dogs, Their Prevalence and Significance of Public health in Sivas). *T Parazit Derg* **21**: 305-309
- BALBO T, ABATE O 1972: Histochemical differentiation of microfilariae of *Dirofilaria immitis*, *Dirofilaria repens* and *Dipetalonema* sp. *Parassitologia* **14**: 239-244
- CANTORAY R, DIK B, GÜLBAHCE S 1990: Konya'da dört köpekte saptanan *Dirofilaria immitis* (Leidy, 1856) olgusu (Quatre cas de *Dirofilaria immitis* (Leidy, 1856) chez les chiens de Konya). *Veterinarium* **1**: 32-34
- CHALIFOUX L, HUNT RD 1971: Histochemical differentiation of *Dirofilaria immitis* and *Dipetalonema reconditum*. *J Am Vet Med Assoc* **158**: 601-605
- CHENG TC 1986: *General Parasitology*. 2th ed. Academic Press, Florida, pp. 544-547
- COŞKUN SZ, TINAR R, AKYOL ÇV 1992: Dogal enfekte köpeklerde *Dirofilaria immitis* mikrofilarlerine ivermektinin etkisi (Efficacy of Ivermectin Against *Dirofilaria immitis* Microfilariae in Naturally Infected dogs). *ÜÜ Vet Fak Derg* **11**: 121-128
- DOGANAY A 1983: Ankara köpeklerinde görülen helmint türleri, bunların yayılışı ve halk sağlığı yönünden önemi (Prevalence of Helminths in Ankara dogs and their potential public health significance). *AÜ Vet Fak Derg* **30**(4): 550-561
- ERDİL, N 1966: Köpeklerde *Dirofilaria immitis* ve *Dipetalonema reconditum* üzerinde araştırma (A Study on *Dirofilaria immitis* and *Dipetalonema reconditum* in dogs). K.K.K. As. Vet. Akademisi, İç Hastalıklar Sube ve Kliniği Çalışmaları, Ankara
- KELLY JD 1979: Canine heartworm disease. *Current Veterinary Therapy VII*. W.B. Saunders, 326-335 pp.
- LINDSEY JR 1965: Identification of canine microfilariae. *J Am Vet Med Assoc* **146**: 1106-1114
- ORTEGA – MORA LM, GOMEZ BAUTISTA M, ROJO – VAZQUEZ FA 1989: The acid phosphatase activity and morphological characteristics of *Dipetalonema dracunculoides* (Cobbold, 1870) microfilariae. *Vet Parasitol* **33**: 187-190
- ROSSIL, ABATE O 1990: Acid phosphatase activity in microfilariae of *Setaria labiati* – papillosa and comparison with other blood microfilariae of dog and horse origin. *Parassitologia* **32**: 339-342
- SARNIÇ H., ALKAN M. 1986: Köpeklerde *Dirofilaria immitis* olguları ve insan sağlığı yönünden önemi (*Dirofilaria immitis* in dogs and its importance for public health). *T Parazit Derg* **1-2**: 169-174
- SCHREY C.F. 1996: Epidemiologische Fallanalyse und klinik der kardiovaskulären *Dirofilariose* (Herzwerkmerkrankung) bei Hunden in Deutschland. PhD Thesis, Freie Universität Berlin, Berlin, Journal Nr. 1942, 125 pp.
- SLOSS MW, KEMP RL, ZAJAC AM 1994.: *Veterinary Clinical Parasitology* 6th ed. Iowa State University Press, Iowa, pp. 107-108
- SOULSBY E.J.L 1982: *Helminths, Arthropods and Protozoa of Domesticated Animals* 7th ed. Bailliere Tindall, London, pp. 307-320
- SAHIN I, EKINCI N, SEN I, ÖZCAN M, GÖDEKMERDAN A 1993: Kayseri yöresi köpeklerinde *Echinococcus granulosus* (Batsch, 1786) ve diğer parazitlerin yayılışı (The distribution and prevalence of *Echinococcus granulosus* (Batsch, 1786) and other parasites in dogs in Kayseri region). *T Parazit Derg* **17**: 69-76
- TASAN E 1983: Elazığ ve yöresindeki köpeklerde filariaların yayılışı (The distribution of Filariasis in Elazığ and its vicinity). *Doga Bilim Derg* **7**: 63-70
- TASAN E 1984: Elazığ kırsal yöre köpeklerinde helmintlerin yayılışı ve insan sağlığı yönünden önemi (The distribution and public health significance of dog helminths in rural districts of Elazığ). *Doga Bilim Derg*. **8**: 160-167
- TINAR R, COŞKUN SZ, DOĞAN H, DEMİR S, AKYOL ÇV, AYDIN L 1989: Bursa yöresi köpeklerinde görülen helmint türleri ve bunların yayılışı (Prevalence of helminths in Bursa stray dogs). *T Parazit Derg* **13**: 113-120
- WHITLOCK HV, PORTER CJ, KELLY JD 1978: The PKW acid phosphatase modification for the recovery histochemical identification of microfilariae of *Dirofilaria immitis* in blood. *Aust Vet Practitioner* **8**: 201-207
- ZEYBEK H 1989: Ankara yöresi köpeklerinde *Dirofilaria immitis* olguları (Les incidences *Dirofilaria immitis* chez les chiens de la region d' Ankara). *Etilik Vet Mikrobiyol Derg* **6**: 1-9
- ZEYBEK H, TATAR N, TOKAY A 1992.: Ankara yöresi kırsal alan köpeklerinde görülen parazitler ve bunların yayılışı (The parasites seen in the rural dogs in Ankara region and their distribution). *Etilik Vet Mikrobiyol Derg* **7**: 17-27



Fig. 1. Pink-stained microfilaria *Dipetalonema reconditum* with naphthol AS-TR phosphatase technique